Take the test. Beat the pest.

Michigan Soybean Cyst Nematode

FREE ANALYSIS

For Free Analysis MUST Have Complete Farmer Name and Address



SCN SAMPLING INSTRUCTIONS:

- A: Always place soil and root samples for SCN in the plastic bag provided and keep cool.
- B: During the growing season, collect samples from margins of diseased areas and inspect roots for SCN females.
- C: Following harvest or before planting, collect multiple soil cores (50-100) at random by walking in a Z- or Wshaped pattern. Mix soil in pail and fill bag with about 1 quart.

Ъ

Deliver to your local MSUE Office

East Lansing, MI 48824-6469

Michigan State University

- D: The Nematode Lab at MSU recommends fields are tested for SCN the fall before every soybean crop.
- E: Refer to MSU Ag Facts E-2199 if more detailed information is required on sampling for nematodes.

SAMPLE DELIVERY INSTRUCTIONS

Deliver or mail (in corrugated box) to

578 Wilson Rd., Room #107

Diagnostic Services

SCN sampling is funded by the Michigan Soybean Promotion Committee through the soybean checkoff.





Funded by Soybean Checkoff Dollars

The Soybean Cyst Nematode (SCN) is a major limiting factor in Michigan's soybean production. It's imperative that problem fields are identified for proper management of this important soybean pathogen. Identification of SCN requires inspection of root tissue and submission of soil samples to a Diagnostic Laboratory such as the one at MSU. Cysts are extracted from the soil to estimate the numbers of eggs and juveniles present, so risk to subsequent soybean crops can be assessed. Recommendations for management are derived from this information.

Because sampling is necessary for SCN identification, the Michigan Soybean Promotion Committee will pay the analysis costs of samples submitted to the MSU program. Please fill out the form completely (one per sample) and either deliver or mail the samples to Diagnostic Services, 578 Wilson Rd., Michigan State University, East Lansing, MI 48824-6469 or deliver the sample to your local MSUE office. Sample results will be returned as quickly as possible. Details for nematode sample collection and care are outlined in MSU Ag Facts Bulletins E-2199 and E-2200 and also on the back of this flyer.

*The normal \$40 SCN Type Test fee will be paid for with MSPC checkoff dollars.

Name		Number of soybean crops grown in	
Address		this field in last 20 years?	
City	Zip	Have SCN resistant varieties been grown in this field? YES NO	
County			
Phone		Circle the SCN source of resistance	
		 in the last variety you planted. 	
Field I.D.	No. Acres	PI 88788 Peking Cyst X Not Sure	

GROWER INFORMATION

* If > 2,500 eggs are found in this sample, would you like an SCN Type Test? YES NO (Requires 45-90 days to complete)

SAMPLE RESULTS do not write below dotted line: lab use only

Risk3

MSU Case #
MSPC #
Date Rec'd

Diagnosis and Recommendations:

Soybean	Cysts			
Cyst	Eggs	J_{2s}		
	J _{2s}			
Lesion				
Root-Knot				
Lance				
Dagger				
Stunt				
Pin				
Spiral				
1. Number per 100cm ³ soil				

Present Crop

Yield of last soybean crop (bu/acre)

Roots²

2. Number per 1.0g root tissue 3. Risk ratings: 0 = none; 1 = low; 2 = moderate; 3 = high

Soil1

Nematodes